

## Recalibrating the Barometer: Echocardiography in Diastolic Dysfunction and the Era of New Algorithms

Maria Estefania Bosco Otto,<sup>1,2</sup> Jorge Eduardo Assef,<sup>3</sup> Gustavo Nishida<sup>3</sup>

Universidade de Brasília,<sup>1</sup> Brasília, DF – Brazil

Hospital DF Star,<sup>2</sup> Brasília, DF – Brazil

Instituto Dante Pazzanese de Cardiologia,<sup>3</sup> São Paulo, SP – Brazil

Diastolic dysfunction (DD) remains a diagnostic challenge, not for lack of available parameters, but because uncertainty emerges when complex physiology is reduced to static labels in echocardiographic reports. In clinical practice, physicians are often less concerned with the specific grade of DD and more interested in its prognostic implications and in whether increased filling pressures may explain dyspnea, guide further investigation, and support therapeutic decisions. It is precisely in this variable, mean left atrial pressure (MLAP) and left ventricular (LV) filling pressure (LVFP), that echocardiography must be most pragmatic, minimizing the number of “indeterminate” reports and offering an operational conclusion grounded in integrated physiology.<sup>1-4</sup>

The structured reasoning that supports this goal began with the 2009 American Society of Echocardiography/European Association of Echocardiography guideline on the evaluation of LV diastolic function,<sup>1</sup> a landmark document that organized modern thinking on diastole, established a shared language for key pathophysiological mechanisms, and systematized the interpretation of echocardiographic parameters. Beyond its conceptual value, it reinforced a critical principle for echocardiography laboratories: the assessment of diastole must translate into a clinically meaningful message, particularly when heart failure with preserved ejection fraction (HFpEF) is suspected.

The 2016 update of the guideline represented another important step by simplifying and improving the reproducibility of multiparametric assessment, focusing on widely available variables suitable for routine use.<sup>2</sup> In practice, however, the post-2016 experience revealed a persistent issue: in the real world, especially among patients with HFpEF, a substantial proportion of studies continued to be classified as having “indeterminate” diastolic function. While methodologically honest, this result is often clinically insufficient.<sup>2-4</sup>

The study by Lababidi et al.<sup>3</sup> represents a pragmatic turning point. In a multicenter cohort validated against invasive hemodynamics, they proposed a stepwise echocardiographic

algorithm to estimate LVFP. The first stage relies on highly feasible measurements, while the second stage resolves discordance or incomplete data using additional parameters supported by strong pathophysiological rationale. Its editorial relevance is direct: the algorithm was designed to reduce the proportion of “indeterminate” cases and improve diagnostic accuracy (from 80% to 86% in patients with HFpEF) in determining LVFP.<sup>3</sup>

Following the publication of Lababidi et al.,<sup>3</sup> the 2025 guideline “recalibrates the barometer” by incorporating these proposed algorithms into a structured strategy for diastolic evaluation and HFpEF diagnosis.<sup>4</sup> This update not only refines the accuracy of DD classification but also improves the identification of increased LVFP, significantly reducing the proportion of cases classified as indeterminate.<sup>4</sup> These algorithms apply to patients in sinus rhythm and without mitral valve conditions that distort the assessment of relaxation and LVFP, such as mitral stenosis of any degree and moderate or severe mitral regurgitation or mitral annular calcification.

However, the 2025 guideline presents two key figures (Figures 2 and 3 of that document<sup>4</sup>) that, at a superficial reading, may appear to compete with one another and therefore generate operational uncertainty. In general terms, Figure 2 of that guideline evaluates the presence of DD, whereas Figure 3 classifies findings based on the estimated MLAP.<sup>4</sup> When interpreted as alternative pathways rather than complementary steps, they may perpetuate uncertainty and lead to conflicting results.

In this context, the letter by Assef & Nishida<sup>5</sup> gains practical relevance. They propose integrating these figures sequentially, redefining Figure 2 as Algorithm 1 (steps for diagnosing DD) and Figure 3 as Algorithm 2 (grading DD and estimating MLAP), both presented in Figure 1 of this Editorial, with clearer transition rules and tie-breaking criteria.<sup>5</sup> This integrated use facilitates interpretation and reduces the likelihood of misclassification. The editorial response from the guideline authors, by clarifying issues related to interpretation and applicability, further contributes to harmonizing understanding and reducing variability in the use of these tools.<sup>6</sup>

In practice, this integration has two immediate consequences. First, it preserves a core set of widely feasible measurements (transmitral Doppler,  $e'$  velocities,  $E/e'$  ratio, and pulmonary pressure estimation) as the initial decision-making step, while directing the use of additional parameters with strong pathophysiological rationale when needed. Second, so-called advanced parameters are placed in their proper role: not as technological embellishments, but as tools to resolve discordance and bring the report closer to the central clinical question, increased versus non-increased LVFP.<sup>4,5</sup>

### Keywords

Left Ventricular Dysfunction; Diastolic Heart Failure; Echocardiography.

**Mailing Address:** Maria Estefania Bosco Otto •

Universidade de Brasília. SQSW 301 Bloco F, Apto 508. Postal code: 70910-900. Brasília, DF – Brazil


E-mail: mariaestefaniaotto@gmail.com

Manuscript received January 12, 2026; revised January 29, 2026; accepted January 29, 2026

Editor responsible for the review: Marcelo Tavares

**DOI:** <https://doi.org/10.36660/abcimg.20260004>

### Algorithm 1 – Steps for diagnosing DD




\*Age-adjusted cutoff values may be considered to identify abnormalities in e' velocity or a reduced E/A ratio.


\*\*After excluding LA enlargement in athletes, anemia, atrial fibrillation, atrial flutter, and mitral valve disease.

$\Delta$ Pre-capillary pulmonary hypertension must be ruled out.

$\ddagger$ After excluding increased LV mass in athletes.



### Algorithm 2 – DD grading & LAP estimation



**Figure 1** – Integrated algorithm for DD (adapted from Assef & Nishida<sup>5</sup> with permission). DD: diastolic dysfunction; IVRT: isovolumic relaxation time; LA: left atrium; LAP: LA pressure; LASr: LA strain, reservoir phase; LAVi: LA volume index; LV: left ventricle; LVMi: LV mass index; S/D: systolic-to-diastolic; PASP: pulmonary artery systolic pressure; TR: tricuspid regurgitation.

Beyond workflow reorganization, the 2025 guideline incorporates two messages that directly reflect daily clinical practice. The first is the recognition of left atrial reservoir strain as an additional variable for diastolic assessment, particularly in populations with preserved ejection fraction, in whom the range of normality is broad and load dependence requires greater interpretative sophistication.<sup>4,7</sup> The second is the reinforcement that mitral annular tissue Doppler  $e'$  velocities should be interpreted in the context of age, acknowledging the physiological decline in relaxation associated with aging.<sup>4</sup> Together, these incorporations move in the same direction: reducing false conflicts among variables and increasing the likelihood of a coherent conclusion when the clinical picture is suggestive.<sup>4,7</sup>

Furthermore, the renewed role of pulmonary vein Doppler is consistent with the historical evolution of the field. As early as 2009, this parameter played a relevant role in inferring LVFP and distinguishing filling patterns.<sup>1</sup> By repositioning it as a tie-breaking variable, contemporary algorithms restore its ability to appropriately reduce the diagnostic gray zone.<sup>1,5</sup>

Integration with clinical algorithms represents the next step in reducing attribution errors in patients with multifactorial dyspnea. Approaches such as  $H_2$ FPEF help estimate pretest probability and identify those who may benefit from additional investigation.<sup>8</sup> Complementarily, the HFA-PEFF algorithm

organizes diagnostic probability and guides decisions regarding functional testing or invasive hemodynamic assessment.<sup>9</sup>

Finally, improved diagnostic precision has an unavoidable practical consequence: HFpEF now has therapies that modify outcomes. Without turning this Editorial into a therapeutic review, it is important to recognize that trials such as EMPEROR-Preserved (empagliflozin) and DELIVER (dapagliflozin) established SGLT2 inhibitors as beneficial interventions in patients with HFpEF.<sup>10,11</sup> The more consistent the diagnosis of an HFpEF phenotype, the more appropriate the application of evidence-based treatment strategies.

In summary, “recalibrating the barometer” in DD means recovering what the 2009 guideline<sup>1</sup> conceptually organized, recognizing what the 2016 guideline<sup>2</sup> simplified, and applying what the 2025 guideline<sup>4</sup> operationalized: a stepwise, integrated, purpose-driven approach to classifying diastolic function and estimating MLAP.<sup>3-7</sup> New diastolic algorithms enable a clinically actionable conclusion, anchored in physiology, regarding the likelihood of increased LVFP. By sequentially integrating diastolic function classification with LVFP estimation, as proposed by Assef & Nishida,<sup>5</sup> inconsistencies observed in recent guidelines are substantially reduced, and echocardiographic reporting gains greater coherence and robustness, particularly when interpreted within the patient’s clinical context.

## References

1. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography. *J Am Soc Echocardiogr.* 2009;22(2):107-33. doi: 10.1016/j.echo.2008.11.023.
2. Nagueh SF, Smiseth OA, Appleton CP, Byrd BF 3rd, Dokainish H, Edvardsen T, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr.* 2016;29(4):277-314. doi: 10.1016/j.echo.2016.01.011.
3. Lababidi H, Rahi W, Smiseth OA, Billick K, Inoue K, Khan FH, et al. New Algorithm for Estimating Left Ventricular Filling Pressure by Echocardiography. *Circulation.* 2025;152(7):424-35. doi: 10.1161/CIRCULATIONAHA.125.074974.
4. Nagueh SF, Sanborn DY, Oh JK, Anderson B, Billick K, Derumeaux G, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography and for Heart Failure with Preserved Ejection Fraction Diagnosis: An Update from the American Society of Echocardiography. *J Am Soc Echocardiogr.* 2025;38(7):537-69. doi: 10.1016/j.echo.2025.03.011.
5. Assef JE, Nishida C. Systematizing Diastolic Function Evaluation: From Algorithms to Practical Approach-Are we Finally Reaching Diagnostic Consistency? *J Am Soc Echocardiogr.* 2026;39(2):235-36. doi: 10.1016/j.echo.2025.09.023.
6. Nagueh SF, Sanborn DY, Oh JK, Anderson B, Billick K, Derumeaux G, et al. Reply to Multiple Letters regarding the American Society of Echocardiography’s Recommendations for the Evaluation of Left Ventricular Diastolic Function. *J Am Soc Echocardiogr.* 2026;39(2):239-40. doi: 10.1016/j.echo.2025.10.010.
7. Nagueh SF, Khan SU. Left Atrial Strain for Assessment of Left Ventricular Diastolic Function: Focus on Populations with Normal LVEF. *JACC Cardiovasc Imaging.* 2023;16(5):691-707. doi: 10.1016/j.jcmg.2022.10.011.
8. Reddy YNV, Carter RE, Obokata M, Redfield MM, Borlaug BA. A Simple, Evidence-Based Approach to Help Guide Diagnosis of Heart Failure with Preserved Ejection Fraction. *Circulation.* 2018;138(9):861-70. doi: 10.1161/CIRCULATIONAHA.118.034646.
9. Pieske B, Tschöpe C, de Boer RA, Fraser AG, Anker SD, Donal E, et al. How to Diagnose Heart Failure with Preserved Ejection Fraction: the HFA-PEFF Diagnostic Algorithm: A Consensus Recommendation from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur Heart J.* 2019;40(40):3297-317. doi: 10.1093/eurheartj/ehz641.
10. Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Böhm M, et al. Empagliflozin in Heart Failure with a Preserved Ejection Fraction. *N Engl J Med.* 2021;385(16):1451-61. doi: 10.1056/NEJMoa2107038.
11. Solomon SD, McMurray JVV, Claggett B, de Boer RA, DeMets D, Hernandez AF, et al. Dapagliflozin in Heart Failure with Mildly Reduced or Preserved Ejection Fraction. *N Engl J Med.* 2022;387(12):1089-98. doi: 10.1056/NEJMoa2206286.



This is an open-access article distributed under the terms of the Creative Commons Attribution License